



Gramin Krishi Mausam Sewa
Agromet Advisory Bulletin for KISHANGANJ District
Bihar Agricultural University
Sabour, Bihar



Agromet Advisory Bulletin

Date : 27-09-2024

Weather Forecast of KISHANGANJ(Bihar) Issued On : 2024-09-27(Valid Till 08:30 IST of the next 5 days)

Parameter	2024-09-28	2024-09-29	2024-09-30	2024-10-01	2024-10-02
Rainfall(mm)	50.0	0.0	0.0	10.0	30.0
Tmax(°C)	24.0	25.0	27.0	27.0	26.0
Tmin(°C)	22.0	24.0	25.0	25.0	24.0
RH-I(%)	95	70	70	80	90
RH-II(%)	65	40	40	50	60
Wind Speed(kmph)	15	10	5	10	15
Wind Direction(Degree)	70	70	100	70	70
Cloud cover(octa)	8	5	5	8	8

Weather Summary/Alert:

According to the medium range forecast issued by the India Meteorological Department for the next five days, the maximum temperature is expected to be between 25.0 °C and 28.0°C . The minimum temperature is expected to be between 23.0°C and 25.0°C . The relative humidity is expected to be between 70 and 95 percent in the morning and 40 to 60 percent in the afternoon and the wind speed is expected to be between 5. 0 km ph. to 20 . 0 km ph. The wind will mostly blow in the South East to west , direction. In the coming days, there is a possibility of light to medium, rainfall along with cloudy weather.

General Advisory:

In the coming days, the weather will be mostly dry and the sky is likely to be light to moderately cloudy, but there is a possibility of medium to light rain on 2 nd October. Keeping in view the coming days, farmer brothers should manage the crops properly. do it, Do not spray crops at this time, spray only when the weather is dry. no water in vegetable beds Let it happen And make proper arrangements for drainage of water from the beds.

SMS Advisory:

In the Coming days till 2nd October light to medium rain fall.

Crop Specific Advisory:

Crop	Crop Specific Advisory
RICE	Crop Specific Advisory RICE Maintain up to 5 cm of water in the paddy field at the time of ear emergence. And at this time, there may be a problem of stem borer insect in paddy. Farmer brothers should keep monitoring their fields regularly. If dry (white) ears are seen if yellow patches (blast disease) or symptoms of stem rot (sheath blight) are seen in the

Crop	Crop Specific Advisory
	paddy ear, then for its protection, spray Propiconazole should be sprayed by making a solution of 1 ml per liter of water.
FINGER MILLET	If symptoms of leaf blast appear on ragi , spraying of tricyclazole @ 1.0 -1.5 ml/liter of water is beneficial. For control of stem fly, spray a solution of Thiomethoxam + Lambdacyhalothrin @ 1 ml/litre of water.
RAPESEED	Those farmer brothers who have not sown rapeseed yet should sow early maturing varieties as soon as possible. Sow rapeseed at 30x15 cm spacing and in 3-4 cm deep furrows.